

ASBESTOS (CHRYSOTILE, AMOSITE, CROCIDOLITE, TREMOLITE, ACTINOLITE, AND ANTHOPHYLLITE)

Asbestos was considered by previous IARC Working Groups in 1972, 1976, and 1987 ([IARC, 1973, 1977, 1987a](#)). Since that time, new data have become available, these have been incorporated in the *Monograph*, and taken into consideration in the present evaluation.

1. Exposure Data

1.1 Identification of the agent

Asbestos is the generic commercial designation for a group of naturally occurring mineral silicate fibres of the serpentine and amphibole series. These include the serpentine mineral chrysotile (also known as ‘white asbestos’), and the five amphibole minerals – actinolite, amosite (also known as ‘brown asbestos’), anthophyllite, crocidolite (also known as ‘blue asbestos’), and tremolite ([IARC, 1973](#); [USGS, 2001](#)). The conclusions reached in this *Monograph* about asbestos and its carcinogenic risks apply to these six types of fibres wherever they are found, and that includes talc containing asbestiform fibres. Erionite (fibrous aluminosilicate) is evaluated in a separate *Monograph* in this volume.

Common names, Chemical Abstracts Service (CAS) Registry numbers and idealized chemical formulae for the six fibrous silicates designated as ‘asbestos’ are presented in [Table 1.1](#). Specific

chemical and physical properties are also presented.

1.2 Chemical and physical properties of the agent

The silicate tetrahedron (SiO_4) is the basic chemical unit of all silicate minerals. The number of tetrahedra in the crystal structure and how they are arranged determine how a silicate mineral is classified.

Serpentine silicates are classified as ‘sheet silicates’ because the tetrahedra are arranged to form sheets. Amphibole silicates are classified as ‘chain silicates’ because the tetrahedra are arranged to form a double chain of two rows aligned side by side. Magnesium is coordinated with the oxygen atom in serpentine silicates. In amphibole silicates, cationic elements such as aluminium, calcium, iron, magnesium, potassium, and sodium are attached to the tetrahedra. Amphiboles are distinguished from one another by their chemical composition. The chemical formulas of asbestos minerals are idealized. In

Table 1.1 Common names, CAS numbers, synonyms, non-asbestos mineral analogues, idealized chemical formulae, selected physical and chemical properties of asbestos minerals

Common Name	CAS No.	Synonyms	Non-Asbestos Mineral Analogue	Idealized Chemical Formula	Colour	Decomposition Temperature (°C)	Other Properties
Asbestos	1332-21-4*	Unspecified		Unspecified			
<i>Serpentine group of minerals</i>							
Chrysotile	12001-29-5*	Serpentine asbestos; white asbestos	Lizardite, antigorite	$[\text{Mg}_3\text{Si}_2\text{O}_5(\text{OH})_4]_n$	White, grey, green, yellowish	600–850	Curled sheet silicate, hollow central core; fibre bundle lengths = several mm to more than 10 cm; fibres more flexible than amphiboles; net positive surface charge; forms a stable suspension in water; fibres degrade in dilute acids
<i>Amphibole group of minerals</i>							
Crocidolite	12001-28-4*	Blue asbestos	Riebeckite	$[\text{NaFe}^{2+}_3\text{Fe}^{3+}_2\text{Si}_8\text{O}_{22}(\text{OH})_2]_n$	Lavender, blue green	400–900	Double chain silicate; shorter, thinner fibres than other amphiboles, but not as thin as chrysotile; fibre flexibility: fair to good; spinnability: fair; resistance to acids: good; less heat resistance than other asbestos fibres; usually contains organic impurities, including low levels of PAHs; negative surface charge in water
Amosite	12172-73-5*	Brown asbestos	Grunerite	$[(\text{Mg}, \text{Fe}^{2+})_7\text{Si}_8\text{O}_{22}(\text{OH})_2]_n$	Brown, grey, greenish	600–900	Double chain silicate; long, straight, coarse fibres; fibre flexibility: somewhat; resistance to acids: somewhat; occurs with more iron than magnesium; negative surface charge in water
Anthophyllite	17068-78-9*	Ferroanthophyllite; azbolen asbestos	Anthophyllite	$[(\text{Mg}, \text{Fe}^{2+})_7\text{Si}_8\text{O}_{22}(\text{OH})_2]_n$	Grey, white, brown-grey, green	NR	Double chain silicate; short, very brittle fibres; resistance to acids: very; relatively rare; occasionally occurs as contaminant in talc deposits ; negative surface charge in water
Actinolite	12172-67-7*	Unspecified	Actinolite	$[\text{Ca}_2(\text{Mg}, \text{Fe}^{2+})_5\text{Si}_8\text{O}_{22}(\text{OH})_2]_n$	Green	NR	Double chain silicate; brittle fibres; resistance to acids: none; occurs in asbestiform and non-asbestiform habit; iron-substituted derivative of tremolite ; common contaminant in amosite deposits; negative surface charge in water
Tremolite	14567-73-8*	Silicic acid; calcium magnesium salt (8:4)	Tremolite	$[\text{Ca}_2\text{Mg}_5\text{Si}_8\text{O}_{22}(\text{OH})_2]_n$	White to pale green	950–1040	Double chain silicate; brittle fibres; acid resistant; occurs in asbestiform and non-asbestiform habit; common contaminant in chrysotile and talc deposits ; negative surface charge in water

* identified as asbestos by CAS Registry

NR, not reported

From [ATSDR \(2001\)](#), [USGS \(2001\)](#), [HSE \(2005\)](#), [NTP \(2005\)](#)

to be more potent than short and thick fibres in the induction of lung cancer in humans. Unfortunately until recently, all of the epidemiological studies that have been conducted used methods for exposure assessment that did not include a determination of fibre size, and thus this issue could not be directly addressed with these studies. As described above, the meta-analysis conducted by [Berman & Crump \(2008a\)](#) considered the effect of fibre size on lung cancer risk by using data from other studies conducted in similar circumstances as the cohort studies. Their analysis did not reveal strong evidence that lung cancer potency was dependent on fibre size. There was weak evidence that long fibres (length > 10 µm) were more potent than short fibres (5 µm < length < 10 µm) in models using all widths ($P = 0.07$). The lack of size-specific data from the studies was a major limitation of this study with regard to estimating size-specific risk estimates. [Stayner et al. \(2008\)](#) published findings from an analysis of the South Carolina asbestos textile cohort in which fibre size specific estimates of lung cancer mortality was evaluated using information from a reanalysis of archived air samples using TEM ([Dement et al., 2008](#)). Long fibres (> 10 µm) and thin fibres (< 0.25 µm) were found to be the strongest predictors of lung cancer mortality in this study.

Another study not part of the prior meta-analyses provides relevant information regarding the question of the relative lung cancer potency of the fibre types. [Loomis et al. \(2009\)](#) carried out a retrospective cohort mortality study of textile workers from four plants in North Carolina that had never been studied before. Workers in this cohort were primarily exposed to chrysotile asbestos that was imported from Quebec. A small amount of amosite was used in an operation in one of the plants. Overall, an excess of lung cancer was observed in this study (SMR, 1.96; 95%CI: 1.73–2.20), which was very similar in magnitude to that reported in the South Carolina cohort study of textile workers ([Hein et al., 2007](#)).

However, the slope for the exposure–response between asbestos exposure and lung cancer was considerably lower than that reported in the South Carolina cohort study. The reasons for these differences in the exposure–response relationships are unknown, but one possible reason may be that quality of the exposure information was superior in the South Carolina study, and that the difference could be explained by an attenuation of the slope due to exposure misclassification in [Loomis et al. \(2009\)](#).

2.2.2 Environmental exposures

Evidence of an association in women between lung cancer and environmental exposures in New Caledonia to field dust containing tremolite and the use of a whitewash (“po”) containing tremolite has been reported ([Luce et al., 2000](#)). A positive association with heavy residential exposure to asbestos was observed in a lung cancer case–control study the Northern Province of South Africa, which is a crocidolite and amosite mining area ([Mzileni et al., 1999](#)). The association was strongest among women who resided in heavily exposed areas (odds ratio [OR], 5.4; 95%CI: 1.3–22.5; $P_{trend} = 0.02$). A study of lung cancer mortality among women in two chrysotile mining regions of Quebec did not result in an increase in lung cancer (SMR, 0.99; 95%CI: 0.78–1.25) relative to women from 60 other areas of Canada ([Camus et al., 1998](#)).

2.2.3 Non-commercial asbestiform amphibole fibres

There is emerging epidemiological evidence that non-commercial amphibole fibres that are asbestiform have carcinogenic potential. These fibres are not technically “asbestos,” and they were never commercially marketed. However, the Working Group felt it was important to discuss the recent evidence concerning these